

Solpadeine PLUS

SOLUBLE TABLETS

Paracetamol, Codeine, Caffeine



- This medicine is for the short term treatment of acute moderate pain when other painkillers have not worked.
- You should only take this product for a maximum of 3 days at a time. If you need to take it for longer than 3 days you should see your doctor or pharmacist for advice.
- This medicine contains codeine which can cause addiction if you take it continuously for more than 3 days. This can give you withdrawal symptoms from the medicine when you stop taking it.
- If you take this medicine for headaches for more than 3 days it can make them worse.

Please read right through this leaflet before you start using this medicine. This medicine is available without prescription, but you still need to use Solpadeine Plus Soluble Tablets carefully to get the best results from them.

- Keep this leaflet you may need to read it again.
- If you have any questions, or if there is anything you do not understand, ask your pharmacist.

In this leaflet:

- What Solpadeine Plus Soluble Tablets do
- Check before you take Solpadeine Plus Soluble Tablets
- How to take Solpadeine Plus Soluble Tablets
- Possible side effects
- How to store Solpadeine Plus Soluble Tablets
- Further information

1. What Solpadeine Plus Soluble Tablets do

Solpadeine Plus Soluble Tablets are for the short term treatment of acute moderate pain which is not relieved by paracetamol, ibuprofen or aspirin alone. They can be used for migraine, headache, dental pain, period pain, strains & sprains, backache, rheumatic pain and sciatica. The tablets contain three active ingredients. Paracetamol and codeine are painkillers. By working on your body in different ways, they combine to relieve pain. Caffeine acts to further help the effectiveness of paracetamol.

2. Check before you take Solpadeine Plus Soluble Tablets

- This medicine contains codeine which can cause addiction if you take it continuously for more than 3 days. This can give you withdrawal symptoms from the medicine when you stop taking it.
- If you take a painkiller for headaches for more than 3 days it can make them worse.



Do not take Solpadeine Plus Soluble Tablets:

- If you have ever had an allergic reaction to paracetamol, caffeine, codeine, other opioid painkillers or to any of the other ingredients (listed in Section 6)
- If you are taking other medicines containing paracetamol or codeine
- If you are under 12 years.



Ask your doctor before you take this medicine:

- If you have liver or kidney disease, including alcoholic liver disease
- If you have bowel problems including blockage of your bowel
- If you have had an operation to remove your gall bladder
- If you have been told by your doctor that you have an intolerance to some sugars
- If you are on a controlled sodium diet. Each tablet dose contains 854 mg of sodium.



If you are taking other medicines

Talk to your doctor or pharmacist before taking these tablets if you are taking any prescribed medicines, particularly metoclopramide or domperidone (for nausea [feeling sick] and vomiting [being sick]), cholestyramine (to lower blood cholesterol); monoamine oxidase inhibitors (MAOIs) prescribed for depression in the last 2 weeks, or substances that make you drowsy (e.g. sedatives, antidepressants or alcohol). If you take blood thinning drugs (anticoagulants e.g. warfarin) and you need to take a pain reliever on a daily basis, talk to your doctor because of the risk of bleeding. But you can still take occasional doses of Solpadeine Plus Soluble Tablets at the same time as anticoagulants.



Pregnancy and breast feeding

Talk to your doctor before taking Solpadeine Plus Soluble Tablets if you are pregnant. Do not use whilst breast feeding unless advised by your doctor.

3. How to take Solpadeine Plus Soluble Tablets



Adults and children aged 12 years and over:
Take 2 tablets dissolved in a full tumbler of water (200ml), every 4 hours as needed.



- Do not take more frequently than every 4 hours.
- Do not take more than 8 tablets in 24 hours.
- Do not take more than the recommended dose.
- Avoid too much caffeine in drinks like coffee and tea. High caffeine intake can cause difficulty sleeping, shaking and an uncomfortable feeling in the chest.
- Do not give to children under 12 years.
- Do not take for more than 3 days. If you need to use this medicine for more than 3 days you must speak to your doctor or pharmacist.

Possible withdrawal effect

This medicine contains codeine and can cause addiction if you take it continuously for more than 3 days. When you stop taking it you may get withdrawal symptoms. You should talk to your doctor or pharmacist if you think you are suffering from withdrawal symptoms.

If you take too many tablets

Immediate medical advice should be sought in the event of an overdose, even if you feel well, because of the risk of delayed, serious liver damage.

If your symptoms continue or your headache becomes persistent, see your doctor.

4. Possible side effects

Some people may have side effects when using this medicine. If you have any unwanted side effects you should seek advice from your doctor, pharmacist or other healthcare professional. Also you can help make sure that medicines remain as safe as possible by reporting any unwanted side effects via the internet at www.mhra.gov.uk/yellowcard; alternatively you can call Freephone 0800 100 3352 (available between 10am-2pm Monday - Friday) or fill in a paper form available from your local pharmacy.

When using this product you may experience:

- Constipation.

Stop taking this medicine and tell your doctor immediately if you experience:

- Severe abdominal pain, dry mouth, nausea and vomiting if you have recently had your gall bladder removed

- Nervousness, dizziness, irritability, drowsiness or restlessness
- Allergic reactions which may be severe such as skin rash and itching sometimes with swelling of the mouth or face or shortness of breath
- Skin rash or peeling or mouth ulcers, itching or sweating
- Breathing problems. These are more likely if you have experienced them before when taking other painkillers such as ibuprofen and aspirin
- Unexplained bruising or bleeding
- Nausea, upset stomach, sudden weight loss, loss of appetite and yellowing of the eyes and skin
- Difficulty in urinating.

If you do get any side effects, even those not mentioned in this leaflet, tell your doctor or pharmacist.



How do I know if I am addicted?

If you take this medicine according to the instructions on the pack, it is unlikely you will become addicted to the medicine. However, if the following apply to you it is important that you talk to your doctor:

- You need to take the medicine for longer periods of time.
- You need to take more than the recommended dose.
- When you stop taking the medicine you feel very unwell but you feel better if you start taking the medicine again.

5. How to store Solpadeine Plus Soluble Tablets

Keep out of the reach and sight of children.
Do not use this medicine after the 'EXP' date shown on the pack.
Store below 30°C.

6. Further information

Active ingredients: Each capsule contains Paracetamol 500 mg, Codeine Phosphate Hemihydrate 8 mg and Caffeine 30 mg.
Other ingredients: Sodium bicarbonate, sorbitol powder (E 420), saccharin sodium, sodium lauryl sulfate, citric acid, sodium carbonate, polyvinylpyrrolidone and silicone fluid.

Packs of Solpadeine Plus Soluble Tablets contain 16 or 32 effervescent tablets.
The marketing authorisation holder is Omega Pharma Ltd., 32 Vauxhall Bridge Road, London, SW1V 2SA, United Kingdom and all enquiries should be sent to this address.

The manufacturer is GlaxoSmithKline Dunganvan Ltd., Co. Waterford, Ireland.
This leaflet was last revised in December 2012.

SOPS01330812v1